

Possible Implications of Small or Large CP Violation in B_d^0 vs $\bar{B}_d^0 \rightarrow J/\psi K_S$ Decays

Zhi-zhong Xing

*Institute of High Energy Physics, P.O. Box 918, Beijing 100039, China **

(Electronic address: xingzz@mail.ihep.ac.cn)

Abstract

We argue that a small or large CP-violating asymmetry $\mathcal{A}_{\psi K_S}$ in B_d^0 vs $\bar{B}_d^0 \rightarrow J/\psi K_S$ decays, which seems to be favored by the recent BaBar or Belle data, might hint at the existence of new physics in B_d^0 - \bar{B}_d^0 mixing. We present a model-independent framework to show how new physics in B_d^0 - \bar{B}_d^0 mixing modifies the standard-model CP-violating asymmetry $\mathcal{A}_{\psi K_S}^{\text{SM}}$. We particularly emphasize that an experimental confirmation of $\mathcal{A}_{\psi K_S} \approx \mathcal{A}_{\psi K_S}^{\text{SM}}$ must not imply the absence of new physics in B_d^0 - \bar{B}_d^0 mixing.

PACS number(s): 11.30.Er, 12.15.Hh, 12.60.Fr, 13.25.Hw

Typeset using REVTeX

*Mailing address

Recently the BaBar and Belle Collaborations have reported their new measurements of the CP-violating asymmetry in B_d^0 vs $\bar{B}_d^0 \rightarrow J/\psi K_S$ decays:

$$\mathcal{A}_{\psi K_S} = \begin{cases} 0.59 \pm 0.14(\text{stat}) \pm 0.05(\text{syst}) , & (\text{BaBar [1]}), \\ 0.99 \pm 0.14(\text{stat}) \pm 0.06(\text{syst}) , & (\text{Belle [2]}). \end{cases} \quad (1)$$

The central values of these two measurements are apparently different from that of the previous CDF measurement, $\mathcal{A}_{\psi K_S} = 0.79 \pm 0.42$ [3]; and they are also different from the result obtained from global analyses of the Cabibbo-Kobayashi-Maskawa (CKM) unitarity triangle in the standard model, $\mathcal{A}_{\psi K_S}^{\text{SM}} = 0.75 \pm 0.06$ [4]. In view of the error bars associated with the BaBar and Belle measurements, it remains too early to claim any serious discrepancy between the experimental result and the standard-model prediction. Nevertheless, one cannot rule out the possibility of $\mathcal{A}_{\psi K_S} < \mathcal{A}_{\psi K_S}^{\text{SM}}$ or $\mathcal{A}_{\psi K_S} > \mathcal{A}_{\psi K_S}^{\text{SM}}$. A small or large CP-violating asymmetry in $B_d \rightarrow J/\psi K_S$ decays should be a clean signal of new physics beyond the standard model.

The purpose of this Brief Report is two-fold. First, we present a model-independent framework to show how new physics in B_d^0 - \bar{B}_d^0 mixing may modify the standard-model quantity $\mathcal{A}_{\psi K_S}^{\text{SM}}$. We find that the possible deviation of $\mathcal{A}_{\psi K_S}$ from $\mathcal{A}_{\psi K_S}^{\text{SM}}$ can fully be described in terms of three independent parameters, including the magnitude and phase of the new-physics contribution to B_d^0 - \bar{B}_d^0 mixing. Second, we point out that the equality $\mathcal{A}_{\psi K_S} = \mathcal{A}_{\psi K_S}^{\text{SM}}$ itself must not mean the absence of new physics in B_d^0 - \bar{B}_d^0 mixing. Indeed there may exist a specific parameter space for the new-physics contribution to B_d^0 - \bar{B}_d^0 mixing, in which the value of $\mathcal{A}_{\psi K_S}$ coincides with that of $\mathcal{A}_{\psi K_S}^{\text{SM}}$. Hence measuring the CP-violating asymmetry $\mathcal{A}_{\psi K_S}$ alone is neither enough to test the standard model nor enough to constrain the possible new physics in B_d^0 - \bar{B}_d^0 mixing.

It is well known that the CP asymmetry $\mathcal{A}_{\psi K_S}$ arises from the interplay of the direct decays of B_d^0 and \bar{B}_d^0 mesons, the B_d^0 - \bar{B}_d^0 mixing in the initial state, and the K^0 - \bar{K}^0 mixing in the final state [5]:

$$\mathcal{A}_{\psi K_S} = -\text{Im} \left(\frac{q}{p} \cdot \frac{V_{cb}V_{cs}^*}{V_{cb}^*V_{cs}} \cdot \frac{q_K^*}{p_K^*} \right), \quad (2)$$

where V_{cb} and V_{cs} are the CKM matrix elements, p and q are the B_d^0 - \bar{B}_d^0 mixing parameters, p_K and q_K are the K^0 - \bar{K}^0 mixing parameters, and the minus sign on the right-hand side of Eq. (2) comes from the CP-odd eigenstate $J/\psi K_S$. In this expression the tiny penguin contributions to the direct transition amplitudes, which may slightly modify the ratio $(V_{cb}V_{cs}^*)/(V_{cb}^*V_{cs})$ [6], have been neglected. Within the standard model $q_K/p_K \approx 1$, $q/p \approx V_{td}/V_{td}^*$ and $(V_{cb}V_{cs}^*)/(V_{cb}^*V_{cs}) \approx 1$ are excellent approximations in the Wolfenstein phase convention for the CKM matrix [7]. Therefore one obtains

$$\mathcal{A}_{\psi K}^{\text{SM}} \approx -\text{Im} \left(\frac{V_{td}}{V_{td}^*} \right) \approx \sin 2\beta, \quad (3)$$

where $\beta \equiv \arg[-(V_{cb}^*V_{cd})/(V_{tb}^*V_{td})] \approx \arg(-V_{td}^*)$ is one of the three inner angles of the CKM unitarity triangle [8]. A recent global analysis of the quark flavor mixing data and the CP-violating observables in the kaon system yields $\sin 2\beta = 0.75 \pm 0.06$ [4].

If the measured value of $\mathcal{A}_{\psi K_S}$ deviates significantly from the standard-model prediction in Eq. (3), it is most likely that the B_d^0 - \bar{B}_d^0 mixing phase q/p consists of unknown new physics contributions. Of course there may also exist new physics in K^0 - \bar{K}^0 mixing, contributing a non-trivial complex phase to $\mathcal{A}_{\psi K_S}$ through q_K/p_K . It is quite unlikely that the tree-level W -mediated decays of B_d^0 and \bar{B}_d^0 mesons are contaminated by any kind of new physics in a significant way [9].

To be specific, we assume that a possible discrepancy between $\mathcal{A}_{\psi K_S}$ and $\mathcal{A}_{\psi K_S}^{\text{SM}}$ mainly results from new physics in B_d^0 - \bar{B}_d^0 mixing. We therefore write down the ratio q/p in terms of the off-diagonal elements of the 2×2 B_d^0 - \bar{B}_d^0 mixing Hamiltonian:

$$\frac{q}{p} = \sqrt{\frac{M_{12}^* - i\Gamma_{12}^*/2}{M_{12} - i\Gamma_{12}/2}} \quad (4)$$

with

$$M_{12} = M_{12}^{\text{SM}} + M_{12}^{\text{NP}} \quad (5)$$

and $\Gamma_{12} = \Gamma_{12}^{\text{SM}}$. Note that $|M_{12}| \gg |\Gamma_{12}|$ is expected to hold both within and beyond the standard model, thus we have $q/p \approx \sqrt{M_{12}^*/M_{12}}$ as a good approximation. The relative magnitude and the phase difference between the new-physics contribution M_{12}^{NP} and the standard-model contribution M_{12}^{SM} are in general unknown. By definition, we may take $|M_{12}| = \Delta M/2$, where $\Delta M = (0.487 \pm 0.014) \text{ ps}^{-1}$ is the experimentally measured mass difference between two mass eigenstates of B_d mesons [10]. Then we parametrize M_{12}^{SM} , M_{12}^{NP} and M_{12} in the following way:

$$\begin{pmatrix} M_{12}^{\text{SM}} \\ M_{12}^{\text{NP}} \\ M_{12} \end{pmatrix} = \begin{pmatrix} R_{\text{SM}} e^{i2\beta} \\ R_{\text{NP}} e^{i2\theta} \\ e^{i2\phi} \end{pmatrix} \frac{\Delta M}{2}, \quad (6)$$

where R_{SM} and R_{NP} are real and positive parameters, θ represents the new-physics phase, and ϕ denotes the effective (overall) phase of B_d^0 - \bar{B}_d^0 mixing. In this case, M_{12}^{SM} , M_{12}^{NP} and M_{12} (or equivalently, $R_{\text{SM}} e^{i2\beta}$, $R_{\text{NP}} e^{i2\theta}$ and $e^{i2\phi}$) form a triangle in the complex plane, as illustrated by Fig. 1. The dual relation between R_{SM} and R_{NP} can be expressed as

$$R_{\text{NP}} = -R_{\text{SM}} \cos 2(\theta - \beta) \pm \sqrt{1 - R_{\text{SM}}^2 \sin^2 2(\theta - \beta)}, \quad (7a)$$

and

$$R_{\text{SM}} = -R_{\text{NP}} \cos 2(\theta - \beta) \pm \sqrt{1 - R_{\text{NP}}^2 \sin^2 2(\theta - \beta)}, \quad (7b)$$

which depends only upon the phase difference $(\theta - \beta)$. There exist two possible solutions for R_{NP} or R_{SM} , corresponding to (\pm) signs on the right-hand side of Eq. (7). Numerically, $R_{\text{SM}} > 0$ and $R_{\text{NP}} \geq 0$ must hold for either solution.

The magnitude of R_{SM} can be calculated in the box-diagram approximation [11]:

$$R_{\text{SM}} = \frac{G_F^2 B_B f_B^2 M_B m_t^2}{6\pi^2 \Delta M} \eta_B F(z) |V_{tb} V_{td}|^2, \quad (8)$$

where G_F is the Fermi constant, B_B is the “bag” parameter describing the uncertainty in evaluation of the hadronic matrix element $\langle B_d^0 | \bar{b} \gamma_\mu (1 - \gamma_5) d | \bar{B}_d^0 \rangle$, M_B is the B_d -meson mass, f_B is the decay constant, m_t is the top-quark mass, η_B denotes the QCD correction factor, V_{tb} and V_{td} are the CKM matrix elements, and $F(z)$ stands for a slowly decreasing monotonic function of $z \equiv m_t^2/M_W^2$ with M_W being the W -boson mass. At present it is difficult to obtain a reliable value for R_{SM} , because quite large uncertainties may arise from the input parameters B_B , f_B and $|V_{td}|$. However, R_{SM} is in general expected to be close to unity, no matter what kind of new physics is hidden in B_d^0 - \bar{B}_d^0 mixing. Note that $R_{\text{SM}} = 1$ must not lead to $R_{\text{NP}} = 0$. There is another solution, $R_{\text{NP}} = -2 \cos 2(\theta - \beta)$ with $\cos 2(\theta - \beta) \leq 0$, corresponding to $R_{\text{SM}} = 1$. On the contrary, $R_{\text{NP}} = 0$ must result in $R_{\text{SM}} = 1$, as indicated by Eq. (7b).

With the help of Eqs. (5) and (6), we recalculate the CP-violating asymmetry $\mathcal{A}_{\psi K_S}$ and arrive at the following result:

$$\mathcal{A}_{\psi K_S} = \sin(2\phi) = R_{\text{SM}} \sin(2\beta) + R_{\text{NP}} \sin(2\theta). \quad (9)$$

Note that R_{NP} , R_{SM} , β , and θ are dependent on one another through Eq. (7). Of course, $|\mathcal{A}_{\psi K}| \leq 1$ holds within the allowed parameter space of R_{NP} and θ . The ratio of $\mathcal{A}_{\psi K_S}$ to $\mathcal{A}_{\psi K_S}^{\text{SM}}$ is given as

$$\xi_{\psi K_S} \equiv \frac{\mathcal{A}_{\psi K_S}}{\mathcal{A}_{\psi K_S}^{\text{SM}}} \approx \frac{\sin(2\phi)}{\sin(2\beta)} = R_{\text{SM}} + R_{\text{NP}} \frac{\sin(2\theta)}{\sin(2\beta)}. \quad (10)$$

In the literature (e.g., Ref. [4]), the value of $\mathcal{A}_{\psi K_S}^{\text{SM}} \approx \sin 2\beta$ is obtained from a global analysis of the experimental data on $|V_{ub}/V_{cb}|$, B_d^0 - \bar{B}_d^0 mixing, B_s^0 - \bar{B}_s^0 mixing, and CP violation in K^0 - \bar{K}^0 mixing. The key assumption in such analyses is that there is no new-physics contribution to the K^0 - \bar{K}^0 , B_d^0 - \bar{B}_d^0 , and B_s^0 - \bar{B}_s^0 mixing systems. If new physics does contribute significantly to the heavy meson-antimeson mixing instead of the light one, one has to discard the direct experimental data on B_d^0 - \bar{B}_d^0 mixing and B_s^0 - \bar{B}_s^0 mixing in analyzing the CKM unitarity triangle. In this case, the resultant constraint on $\sin 2\beta$ becomes somehow looser. One may observe, from the figures of the CKM unitarity triangle in Refs. [4,8], that $0.6 \leq \sin 2\beta \leq 0.8$ is a quite generous range constrained by current data on $|V_{ub}/V_{cb}|$ and CP violation in K^0 - \bar{K}^0 mixing. Given such a generously allowed region for $\mathcal{A}_{\psi K_S}^{\text{SM}}$, we conclude that $\xi_{\psi K_S} > 0$ is definitely assured. Using $\mathcal{A}_{\psi K_S}^{\text{SM}} = 0.75 \pm 0.06$ [4] for illustration, we obtain

$$\xi_{\psi K_S} = \begin{cases} 0.79 \pm 0.26, & (\text{BaBar}), \\ 1.32 \pm 0.30. & (\text{Belle}). \end{cases} \quad (11)$$

We see that the BaBar measurement seems to indicate $\xi_{\psi K_S} < 1$, while the Belle measurement seems to imply $\xi_{\psi K_S} > 1$. If either possibility could finally be confirmed with more precise experimental data from B -meson factories, it would be a very clean signal of new physics [12]. If the further data of both BaBar and Belle Collaborations turn to coincide with each other and lead to $\xi_{\psi K_S} \approx 1$, however, one cannot draw the conclusion that there is no new physics in B_d^0 - \bar{B}_d^0 mixing.

Now let us show why $\mathcal{A}_{\psi K_S} = \mathcal{A}_{\psi K_S}^{\text{SM}}$ must not imply the absence of new physics in $B_d^0\text{-}\bar{B}_d^0$ mixing. Taking $\xi_{\psi K_S} = 1$ and using Eq. (7), we obtain the following equation constraining the allowed values of θ :

$$(1 + R_{\text{SM}}) \tan^2 2\theta - 2R_{\text{SM}} \tan 2\beta \tan 2\theta - (1 - R_{\text{SM}}) \tan^2 2\beta = 0. \quad (12)$$

Then it is straightforward to find out two solutions for $\tan 2\theta$:

$$\tan 2\theta = \tan 2\beta, \quad (13a)$$

or

$$\tan 2\theta = \tan 2\beta \frac{R_{\text{SM}} - 1}{R_{\text{SM}} + 1}. \quad (13b)$$

Note that solution (13a) corresponds to $R_{\text{SM}} + R_{\text{NP}} = 1$. Solution (13b) implies that $|\tan 2\theta| \ll \tan 2\beta$ may hold, if R_{SM} is remarkably close to 1. Although the afore-obtained region of θ is quite specific, it does exist and give rise to $\xi_{\psi K_S} = 1$. Therefore, an experimental confirmation of $\xi_{\psi K_S} \approx 1$ cannot fully rule out the possibility of new physics hidden in $B_d^0\text{-}\bar{B}_d^0$ mixing.

Theoretically, the information on R_{NP} and θ can only be obtained from specific models of new physics (e.g., the supersymmetric extensions of the standard model [12]). An interesting possibility is that the new-physics contribution conserves CP (i.e., $\theta = 0$ [13]) and all observed CP-violating phenomena in weak interactions are attributed to the non-trivial phase in the CKM matrix. In this scenario, we obtain

$$\mathcal{A}_{\psi K_S} = \sin(2\phi) = R_{\text{SM}} \sin(2\beta). \quad (14)$$

Obviously $\mathcal{A}_{\psi K_S}/\mathcal{A}_{\psi K_S}^{\text{SM}} = R_{\text{SM}} \leq 1$ is required, in order to understand the present BaBar and Belle measurements.

It becomes clear that the measurement of $\mathcal{A}_{\psi K_S}$ itself is not enough to test the self-consistency of the standard model or to pin down possible new physics hidden in $B_d^0\text{-}\bar{B}_d^0$ mixing. For either purpose one needs to study the CP-violating asymmetries in some other nonleptonic B -meson decays, although most of them are not so clean as B_d^0 vs $\bar{B}_d^0 \rightarrow J/\psi K_S$ decays in establishing the relations between the CP-violating observables and the fundamental CP-violating parameters [14].

In summary, we have discussed possible implications of a small or large CP-violating asymmetry in B_d^0 vs $\bar{B}_d^0 \rightarrow J/\psi K_S$ decays. While such an effect could be attributed to new physics in $K^0\text{-}\bar{K}^0$ mixing, it is most likely to result from new physics in $B_d^0\text{-}\bar{B}_d^0$ mixing. Model-independently, we have formulated the basic features of new-physics effects on CP violation in $B_d \rightarrow J/\psi K_S$. We have also pointed out that an experimental confirmation of $\mathcal{A}_{\psi K_S} \approx \mathcal{A}_{\psi K_S}^{\text{SM}}$ must not imply the absence of new physics in $B_d^0\text{-}\bar{B}_d^0$ mixing. An extensive study of all hadronic B -meson decays and CP asymmetries is desirable, in order to test the standard model and probe possible new physics at some higher energy scales.

The author would like to thank L. Wolfenstein for numerous useful discussions.

REFERENCES

- [1] BaBar Collaboration, B. Aubert *et al.*, hep-ph/0107013.
- [2] Belle Collaboration, A. Abashian *et al.*, hep-ph/0107061.
- [3] CDF Collaboration, T. Affolder *et al.*, Phys. Rev. D **61**, 072005 (2000).
- [4] F. Caravaglios, F. Parodi, P. Roudeau, and A. Stocchi, hep-ph/0002171; and references therein.
- [5] D. Du, I. Dunietz, and D.D. Wu, Phys. Rev. D **34**, 3414 (1986); hep-ph/9703458.
- [6] See, e.g., Z.Z. Xing, Phys. Lett. B **353**, 313 (1995); D. Du and Z.Z. Xing, Phys. Lett. B **312**, 199 (1993).
- [7] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1984). A general classification of the parametrizations (phase conventions) for the CKM matrix can be found: H. Fritzsch and Z.Z. Xing, Phys. Lett. B **413**, 396 (1997); Phys. Rev. D **57**, 594 (1998); Prog. Part. Nucl. Phys. **45**, 1 (2000).
- [8] Particle Data Group, D.E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [9] See, e.g., Y. Nir, Lectures presented in the 20th SLAC Summer Institute, SLAC-PUB-5874 (1992).
- [10] A. Stocchi, hep-ph/0010222 (talk at the XXXth International Conference on High Energy Physics, July 2000, Osaka, Japan).
- [11] A.I. Sanda and Z.Z. Xing, Phys. Rev. D **56**, 6866 (1997); Z.Z. Xing, Eur. Phys. J. C **4**, 283 (1998).
- [12] See, e.g., A.L. Kagan and M. Neubert, Phys. Lett. B **492**, 115 (2000); J.P. Silva and L. Wolfenstein, Phys. Rev. D **63**, 056001 (2001); G. Eyal, Y. Nir, and G. Perez, JHEP **0008**, 028 (2000); Z.Z. Xing, hep-ph/0008018 (unpublished); A.J. Buras and R. Buras, Phys. Lett. B **501**, 223 (2001); A. Masiero, M. Piai, and O. Vives, Phys. Rev. D **64**, 055008 (2001); H. Fritzsch and Z.Z. Xing, Phys. Lett. B **506**, 109 (2001); R. Fleischer and T. Mannel, Phys. Lett. B **506**, 311 (2001); A. Buras, hep-ph/0101336; A. Ali and E. Lunghi, hep-ph/0105200.
- [13] J.M. Soares and L. Wolfenstein, Phys. Rev. D **47**, 1021 (1993); A.I. Sanda and Z.Z. Xing, in Ref. [11].
- [14] For a review with extensive references, see: *The BABAR Physics Book*, edited by P.F. Harrison and H.R. Quinn, Report No. SLAC-R-504 (1998).

FIGURES




FIG. 1. Triangular relation of M_{12}^{SM} , M_{12}^{NP} and M_{12} (rescaled by $\Delta M/2$) in the complex plane.